

Worksheets to print out from sofatutor.com

Word Problems with Quadratic Equations







The complete package, **including all tasks**, **hints**, **solutions**, **and solution approaches**, is available to all subscribers of sofatutor.com



Name the methods for solving quadratic equations.

Choose the correct methods.

$$ax^2 + bx + c = 0$$

$$x^2 + bx + c = 0$$


We always have to use the quadratic formula to get the correct solutions.
We can draw the parabola corresponding to the quadratic equation. The x -intercepts are the desired solutions.
We can factor the left-hand side of the quadratic equation $x^2+bx+c=0$ and then solve the two resulting linear equations.
The desired solution is always $x=c$
We can complete the quadratic polynomial on the left-hand side to a square and solve the equation by taking the square root.

Our hints for the tasks




Name the methods for solving quadratic equations.

1. Hint



The solutions of the corresponding quadratic equation are $\,x=10\,$ and $\,x=90.$

2. Hint



This is the quadratic formula.

3. Hint

The zero factor property states that if a product is equal to zero, then one of its factors must also equal zero.

4. Hint

It doesn't matter at all which method you choose to find the solutions. If the solutions exist, then they are always the same.



Solutions and solution approaches for the tasks



Name the methods for solving quadratic equations.

Answer key: B, C, E

If you have to solve a quadratic equation like $ax^2 + bx + c = 0$, or with a = 1, $x^2 + bx + c = 0$, which you can derive by dividing the equation $ax^2 + bx + c = 0$ by a, then you can decide from several methods:

- 1. You could factor the left-hand side of the equation $x^2 + bx + c = 0$ to (x+d)(x+e) = 0 and get the solutions x = -d and x = -e.
- 2. You can complete the square to get $(x+e)^2=d$. You can solve this equation by taking the square root of both sides.
- 3. You can use the quadratic formula for solving $ax^2 + bx + c = 0$.
- 4. You also could draw the parabola corresponding to $f(x) = ax^2 + bx + c$. The x-intercepts of this parabola are the desired solutions.

It doesn't matter which method you choose, as the solutions, if they exist, are always the same.

